

Renovate[®] 3

Specialty Herbicide

SPECIMEN



Aquatic Sites: For control of emerged, submersed and floating aquatic plants in aquatic sites such as ponds, lakes, reservoirs, non-irrigation canals, seasonal irrigation waters and ditches which have little or no continuous outflow, marshes and wetlands, including broadleaf and woody vegetation on banks and shores within or adjacent to these and other aquatic sites.

For use in New York State, comply with Section 24(c) Special Local Need labeling for Renovate[®] 3, SLN NY-060001

Active Ingredient

triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid,
triethylamine salt44.4%

Other Ingredients55.6%

TOTAL100.0%

Acid equivalent: triclopyr - 31.8% - 3 lb/gal

Keep Out of Reach of Children

DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

NOTICE: Read the entire label. Use only according to label directions. Before using this product, read *Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies* at end of label booklet. If terms are unacceptable, return at once unopened.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Renovate is a registered trademark of Dow AgroSciences LLC.

Produced for: **SePRO Corporation**
11550 North Meridian Street
Suite 600, Carmel, IN 46032, U.S.A.

EPA Reg. 62719-37-67690
FPL20120201

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

DANGER

Corrosive. Causes Irreversible Eye Damage. Harmful If Swallowed Or Absorbed Through Skin. Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reaction In Some Individuals. Do not get in eyes or on skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Shoes plus socks;
- Protective eyewear; and
- Chemical resistant gloves (≥ 14 mils) such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the WPS

[40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Table of Contents

Page

Precautionary Statements.....	1
Hazard to Humans and Domestic Animals	1
Personal Protective Equipment (PPE).....	1
Engineering Controls	1
User Safety Recommendations	1
First Aid	1
Environmental Hazards	2
Physical or Chemical Hazards.....	2
Directions for Use	2
Agricultural Use Requirements	2
Non-Agricultural Use Requirements	2
Storage and Disposal	6
General Information for Aquatic and Wetland Sites.....	2
General Use Precautions and Restrictions	2
Grazing and Haying Restrictions	2
Avoiding Injurious Spray Drift.....	2
Spray Drift Management.....	3
Aerial Drift Reduction Advisory	3
Plants Controlled	3
Woody Plant Species.....	3
Annual and Perennial Broadleaf Weeds	4
Aquatic Weeds.....	4
Application Methods	4
Floating and Emerged Weeds	4
Submerged Weeds	4
Wetland Sites.....	5
Terrestrial Sites Associated with Wetland Areas.....	5
Terms and Conditions of Use.....	7
Warranty Disclaimer.....	7
Inherent Risks of Use.....	7
Limitation of Remedies	7

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call INFOTRAC at 1-800-535-5053.</p>	
<p>Note to Applicator: Allergic skin reaction is not expected from exposure to spray mixtures of Renovate 3 herbicide when used as directed.</p>	
<p>Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.</p>	

ENVIRONMENTAL HAZARDS

Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Under certain conditions, treatment of aquatic weeds can result in oxygen depletion or loss due to decomposition of dead plants, which may contribute to fish suffocation. This loss can cause fish suffocation. Therefore, to minimize this hazard, do not treat more than one-third to one-half of the water area in a single operation and wait at least 10 to 14 days between treatments. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Consult with the State agency for fish and game before applying to public water to determine if a permit is needed.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store the product near heat or open flame.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all *Directions for Use* carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls;
- Shoes plus socks;
- Protective eyewear; and
- Chemical-resistant gloves (≥ 14 mils) such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: For applications to non-cropland areas, do not allow entry into areas until sprays have dried, unless applicator and other handler PPE is worn.

GENERAL INFORMATION FOR AQUATIC AND WETLAND SITES

Use Renovate 3 herbicide for control of emerged, submersed and floating aquatic plants in aquatic sites such as ponds, lakes, reservoirs, non-irrigation canals, and ditches which have little or no continuous outflow, marshes and wetlands, including broadleaf and woody vegetation on banks and shores within or adjacent to these and other aquatic sites.

Obtain Required Permits: Consult with appropriate state or local water authorities before applying this product to public waters. State or local public agencies may require permits.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

For use in New York State, comply with Section 24(c) Special Local Need labeling for Renovate® 3, SLN NY-060001.

When applying this product in tank mix combination, follow all applicable use directions, precautions and limitations on each manufacturer's label.

Chemigation: Do not apply this product through any type of irrigation system.

Irrigation: Do not use treated water for irrigation for 120 days following application. As an alternative to waiting 120 days, treated water may be used for irrigation once the triclopyr level in the intake water is determined to be non-detectable by laboratory analysis (immunoassay). There is no restriction on use of water from the treatment area to irrigate established grasses.

Water treated with Renovate 3 may not be used for irrigation purposes for 120 days after application or until residue levels of Renovate 3 are determined by laboratory analysis, or other appropriate means of analysis, to be 1 ppb or less.

Seasonal Irrigation Waters: Renovate 3 may be applied during the off-season to surface waters that are used for irrigation on a seasonal basis provided that there is a minimum of 120 days between applying Renovate 3 and the first use of treated water for irrigation purposes, or until residue levels of Renovate 3 are determined by laboratory analysis, or other appropriate means of analysis, to be 1 ppb or less.

Irrigation Canals/Ditches: Do not apply Renovate 3 to irrigation canals/ditches unless the 120-day restriction on irrigation water usage can be observed or residue levels of Renovate 3 are determined by laboratory analysis, or other appropriate means of analysis, to be 1 ppb or less.

Do not apply Renovate 3 directly to, or otherwise permit it to come into direct contact with grapes, tobacco, vegetable crops, flowers, or other desirable broadleaf plants, and do not permit spray mists containing it to drift into them.

- **Do not** apply to salt water bays or estuaries.
- **Do not** apply directly to un-impounded rivers or streams.
- **Do not** apply on ditches or canals currently being used to transport irrigation water or that will be used for irrigation within 4 months following treatment. It is permissible to treat irrigation and non-irrigation ditch banks.
- **Do not** apply where runoff water may flow onto agricultural land as injury to crops may result.
- When making applications to control unwanted plants on banks or shorelines of moving water sites, minimize overspray to open water.
- The use of a mist blower is not recommended.

Grazing and Haying Restrictions

Except for lactating dairy animals, there are no grazing restrictions following application of this product.

- **Grazing Lactating Dairy Animals:** Do not allow lactating dairy animals to graze treated areas until the next growing season following application of this product.
- Do not harvest hay for 14 days after application.
- Grazed areas of non-cropland and forestry sites may be spot treated if they comprise no more than 10% of the total grazable area.

Slaughter Restrictions: During the season of application, withdraw livestock from grazing treated grass at least 3 days before slaughter.

Avoiding Injurious Spray Drift

Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. It is suggested that a continuous smoke column at or near the spray site or a smoke generator on the spray equipment be used to detect air movement, lapse conditions, or temperature inversions (stable air). If the smoke layers or indicates a potential of hazardous spray drift, do not spray.

Aerial Application: For aerial application near susceptible crops, apply through a Microfoil† or Thru-Valve boom†, or use a drift control additive labeled for aquatic use. Other drift reducing systems or thickened sprays prepared by using high viscosity inverting systems may be used if they are made as drift-free mixtures containing thickening agents labeled for use in aquatics or applications made with the Microfoil or Thru-Valve boom. Keep spray pressures low enough to provide coarse spray droplets. Spray boom should be no longer than 3/4 of the rotor length. Do not use a thickening

agent with the Microfoil or Thru-Valve booms, or other systems that cannot accommodate thick sprays. Spray only when the wind velocity is low (follow state regulations). Avoid application during air inversions. If a spray thickening agent is used, follow all use recommendations and precautions on the product label.

†Reference within this label to a particular piece of equipment produced by or available from other parties is provided without consideration for use by the reader at its discretion and subject to the reader's independent circumstances, evaluation, and expertise. Such reference by SePRO Corporation is not intended as an endorsement of such equipment, shall not constitute a warranty (express or implied) of such equipment, and is not intended to imply that other equipment is not available and equally suitable. Any discussion of methods of use of such equipment does not imply that the reader should use the equipment other than is advised in directions available from the equipment's manufacturer. The reader is responsible for exercising its own judgment and expertise, or consulting with sources other than SePRO Corporation, in selecting and determining how to use its equipment.

Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications:

1. The distance of the outer most operating nozzles on the boom must not exceed 3/4 the length of the rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following *Aerial Drift Reduction Advisory*. [This information is advisory in nature and does not supersede mandatory label requirements.]

Aerial Drift Reduction Advisory

Information on Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see *Wind, Temperature and Humidity, and Temperature Inversions*).

Controlling Droplet Size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length: For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind: Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential.

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Applications should not occur during a local, low level temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of the smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Ground Equipment: To aid in reducing spray drift, Renovate 3 should be used in thickened (high viscosity) spray mixtures using a labeled drift control additive, high viscosity invert system, or equivalent as directed by the manufacturer. With ground equipment, spray drift can be reduced by keeping the spray boom as low as possible; by applying 20 gallons or more of spray per acre; by keeping the operating spray pressures at the lower end of the manufacturer's recommended pressures for the specific nozzle type used (low pressure nozzles are available from spray equipment manufacturers); and by spraying when wind velocity is low (follow state regulations). In hand-gun applications, select the minimum spray pressure that will provide adequate plant coverage (without forming a mist). Do not apply with nozzles that produce a fine-droplet spray.

High Volume Leaf-Stem Treatment: To minimize spray drift, do not use pressure exceeding 50 psi at the spray nozzle and keep sprays no higher than brush tops. A labeled thickening agent may be used to reduce drift.

Plants Controlled

Woody Plant Species

alder	choke cherry
arrowwood	cottonwood
ash	crataegus (hawthorn)
aspen	locust
bear clover (bearmat)	maleleuca (seedlings)
beech	maples
birch	mulberry
blackberry	oaks
blackgum	poison ivy
Brazilian pepper	poison oak
casacara	poplar
ceanothus	salt-bush (<i>Baccharis</i> spp.)
cherry	sweetgum
Chinese tallow	waxmyrtle
chinquapin	willow

Annual and Perennial Broadleaf Weeds

burdock	smart weed
Canada thistle	tansy ragwort
curly dock	tropical sodaapple
elephant ear	vetch
plantain	wild lettuce

Aquatic Weeds

alligatorweed	Pennywort
American lotus	phragmites
American frogbit	pickerelweed
aquatic sodaapple	purple loosestrife
Eurasian watermilfoil	waterhyacinth
milfoil species	waterlily
nuphar (spatterdock)	watershield
parrotfeather†	water primrose

† Retreatment may be needed to achieve desired level of control.

APPLICATION METHODS

FLOATING AND EMERGED WEEDS

For control of waterhyacinth, alligatorweed (see specific directions below), and other susceptible emerged and floating herbaceous weeds and woody plants, apply 1 1/2 to 6 lb ae of triclopyr (2 to 8 quarts of Renovate 3) per acre as a foliar application using surface or aerial equipment. Use higher rates in the rate range when plants are mature, when the weed mass is dense, or for difficult to control species. Repeat as necessary to control regrowth and plants missed in the previous operation, but do not exceed a total of 6 lb ae of triclopyr (8 quarts of Renovate 3) per acre per annual growing season.

Use a non-ionic surfactant in the spray mixture to improve control. Follow all directions and use precautions on the aquatic surfactant label. Apply when plants are actively growing.

Surface Application

Use a spray boom, handgun or other similar suitable equipment mounted on a boat or vehicle. Thorough wetting of foliage is essential for maximum effectiveness. Use 20 to 200 gallons per acre of spray mixture. Special precautions such as the use of low spray pressure, large droplet producing nozzles or addition of a labeled thickening agent may minimize spray drift in areas near sensitive crops.

Aerial Application (Helicopter Only)

Apply with a helicopter using a Microfoil or Thru-Valve boom, or a drift control additive in the spray solution. Apply in a minimum of 10 gallons of total spray mix per acre. Do not apply when weather conditions favor drift to sensitive areas. See label section on aerial application directions and precautions.

Waterhyacinth (*Eichhornia crassipes*)

Apply Renovate 3 at 1 1/2 to 6 lb ae of triclopyr (2 to 8 quarts of Renovate 3) per acre to control waterhyacinth. Apply when plants are actively growing. Use the higher rate in the rate range when the weed mass is dense. It is important to thoroughly wet all foliage with the spray mixture. Use a non-ionic surfactant in the spray mixture. A repeat treatment may be needed to control regrowth or plants missed in the previous treatment.

Alligatorweed (*Alternanthera philoxeroides*)

Apply Renovate 3 at 2 to 6 lb ae of triclopyr (3 to 8 quarts of Renovate 3) per acre to control alligatorweed. It is important to thoroughly wet all foliage with the spray mixture. For best results, add an approved non-ionic aquatic surfactant to the spray mixture. Alligatorweed growing outside the margins of a body of water can be controlled with this treatment. However, alligatorweed growing in water will only be partially controlled. Top growth above the water will be controlled, but the plant will likely regrow from tissue below the water surface.

Precautions for Potable Water Intakes – Lakes, Reservoirs, Ponds:

For applications of Renovate 3 to control floating and emerged weeds in lakes, reservoirs or ponds that contain a functioning potable water intake for human consumption, see chart below to determine the minimum setback distances of the application from the functioning potable water intakes.

Renovate 3 Application Rate				
Area Treated (acres)	2 qt/acre	4 qt/acre	6 qt/acre	8 qt/acre
	Setback Distance (ft)			
<4	0	200	400	500
>4 - 8	0	200	700	900
>8 - 16	0	200	700	1,000
>16	0	200	900	1,300

Note: Existing potable water intakes which are no longer in use, such as those replaced by potable water wells or connections to a municipal water system, are not considered to be functioning potable water intakes. These setback restrictions do not apply to terrestrial applications made adjacent to potable water intakes.

To apply Renovate 3 around and within the distances noted above from a functioning potable water intake, the intake must be turned off until the triclopyr level in the intake water is determined to be 0.4 parts per million (ppm) or less by laboratory analysis or immunoassay.

- **Recreational Use of Water in Treatment Area:** There are no restrictions on use of water in the treatment area for recreational purposes, including swimming and fishing.
- **Livestock Use of Water from Treatment Area:** There are no restrictions on livestock consumption of water from the treatment area.

SUBMERGED WEEDS

For control of Eurasian watermilfoil (*Myriophyllum spicatum*) and other susceptible submerged weeds in ponds, lakes, reservoirs, and in non-irrigation canals or ditches that have little or no continuous outflow, apply Renovate 3 as either a surface or subsurface application. Rates should be selected according to the rate chart below to provide a triclopyr concentration of 0.75 to 2.5 ppm ae in treated water. Use higher rates in the rate range in areas of greater water exchange. These areas may require a repeat application. However, total application of Renovate 3 must not exceed an application rate of 2.5 ppm of triclopyr for the treatment area per annual growing season.

Apply in spring or early summer when Eurasian watermilfoil or other submerged weeds are actively growing.

Areas near susceptible crops or other desirable broadleaf plants may be treated by subsurface injection applied by boat to avoid spray drift.

Subsurface Application

Apply desired amount of Renovate 3 per acre directly into the water through boat-mounted distribution systems. When treating target plants that are 6 feet below the surface of the water, trailing hoses should be used along with an aquatic approved sinking agent (except California).

Surface Application

Apply the desired amount of Renovate 3 as either a concentrate or a spray mixture in water. However, use a minimum spray volume of 5 gallons per acre. Do not apply when weather conditions favor drift to sensitive areas.

Average water depth (feet) x 0.905 x target concentration (ppm) = gallons of Renovate 3 per surface acre treated.

Example: to achieve a 2 ppm concentration of triclopyr in water averaging 4 feet deep

4 x 0.905 x 2 ppm = 7.2 gallons of Renovate 3 per surface acre treated.

Concentration of Triclopyr Acid in Water (ppm ae)					
Water Depth (ft)	0.75 ppm	1.0 ppm	1.5 ppm	2.0 ppm	2.5 ppm
	Gallons of Renovate 3 per Surface Acre at Specified Depth				
1	0.7	0.9	1.4	1.8	2.3
2	1.4	1.8	2.7	3.6	4.6
3	2.1	2.7	4.1	5.4	6.8
4	2.7	3.6	5.4	7.2	9.1
5	3.4	4.5	6.8	9.0	11.3
6	4.1	5.4	8.1	10.9	13.6
7	4.8	6.3	9.5	12.7	15.8
8	5.5	7.2	10.9	14.5	18.1
9	6.1	8.1	12.2	16.3	20.4
10	6.8	9.0	13.6	18.1	22.6
15	10.2	13.6	20.4	27.2	33.9
20	13.6	18.1	27.2	36.2	45.3

Precautions for Potable Water Intakes – Lakes, Reservoirs, Ponds:

For applications of Renovate 3 to control submerged weeds in lakes, reservoirs or ponds that contain a functioning potable water intake for human consumption, see the chart below to determine the minimum setback distances of the application from the functioning potable water intakes.

Concentration of Triclopyr Acid in Water (ppm ae)					
Area Treated (acres)	0.75 ppm	1.0 ppm	1.5 ppm	2.0 ppm	2.5 ppm
	Required Setback Distance (ft) from Potable Water Intake				
<4	300	400	600	800	1,000
>4 - 8	420	560	840	1,120	1,400
>8 - 16	600	800	1,200	1,600	2,000
>16 - 32	780	1,040	1,560	2,080	2,600
>32 acres, calculate a setback using the formula for the appropriate rate	Setback (ft) = $(800 \times \ln(\text{acres}) - 160) / 3.33$	Setback (ft) = $(800 \times \ln(\text{acres}) - 160) / 2.50$	Setback (ft) = $(800 \times \ln(\text{acres}) - 160) / 1.67$	Setback (ft) = $(800 \times \ln(\text{acres}) - 160) / 1.25$	Setback (ft) = $(800 \times \ln(\text{acres}) - 160)$

Example Calculation 1: to apply 2.5 ppm Renovate 3 to 50 acres:

$$\begin{aligned} \text{Setback in feet} &= (800 \times \ln(50 \text{ acres}) - 160) \\ &= (800 \times 3.912) - 160 \\ &= 2,970 \text{ feet} \end{aligned}$$

Example Calculation 2: to apply 0.75 ppm Renovate 3 to 50 acres:

$$\begin{aligned} \text{Setback in feet} &= \frac{(800 \times \ln(50 \text{ acres}) - 160)}{3.33} \\ &= \frac{(800 \times 3.912) - 160}{3.33} \\ &= 892 \text{ feet} \end{aligned}$$

Note: Existing potable water intakes which are no longer in use, such as those replaced by potable water wells or connections to a municipal water system, are not considered to be functioning potable water intakes. These setback restrictions do not apply to terrestrial applications made adjacent to potable water intakes.

To apply Renovate 3 around and within the distances noted above from a functioning potable water intake, the intake must be turned off until the triclopyr level in the intake water is determined to be 0.4 parts per million (ppm) or less by laboratory analysis or immunoassay.

- **Recreational Use of Water in Treatment Area:** There are no restrictions on use of water in the treatment area for recreational purposes, including swimming and fishing.
- **Livestock Use of Water from Treatment Area:** There are no restrictions on livestock consumption of water from the treatment area.

WETLAND SITES

Wetlands include flood plains, deltas, marshes, swamps, bogs, and transitional areas between upland and lowland sites. Wetlands may occur within forests, wildlife habitat restoration and management areas and similar sites as well as areas adjacent to or surrounding domestic water supply reservoirs, lakes and ponds.

For control of woody plants and broadleaf weeds in these sites, follow use directions and application methods on this label for terrestrial sites associated with wetland areas.

Use Precautions: Minimize overspray to open water when treating target vegetation in and around non-flowing, quiescent or transient water. When making applications to control unwanted plants on banks or shorelines of flowing water, minimize overspray to open water. **Note:** Consult local public water control authorities before applying this product in and around public water. Permits may be required to treat such areas.

Purple Loosestrife (*Lythrum salicaria*)

Purple loosestrife can be controlled with foliar applications of Renovate 3. For broadcast applications, use a minimum of 4 1/2 to 6 lb ae of triclopyr (6 to 8 quarts of Renovate 3) per acre. Apply Renovate 3 when purple loosestrife is at the bud to mid-flowering stage of growth. Follow-up applications for control of regrowth should be made the following year in order to achieve increased control of this weed species. For all applications, a non-ionic surfactant labeled for aquatics should be added to the spray mixture. Follow all directions and use precautions on the label of the surfactant. Thorough wetting of the foliage and stems is necessary to achieve satisfactory control. A minimum spray volume of 50 gallons per acre is recommended for ground broadcast applications.

If using a backpack sprayer, a spray mixture containing 1% to 1.5% Renovate 3 or 5 to 7.6 fl oz of Renovate 3 per 4 gallons of water should be used. All purple loosestrife plants should be thoroughly wetted.

Phragmites (*Phragmites australis*)

Phragmites can be selectively controlled with foliar applications of Renovate 3. For broadcast applications, a minimum of 2 1/4 lb ae of triclopyr (3 quarts of Renovate 3) per acre should be used. For optimum control, apply Renovate 3 when phragmites is in the early stage of growth, 1/2 to 3 feet in height, prior to seed head development. Follow-up applications for control of regrowth may be made the following year in order to achieve increased control of this weed species. For all applications, a non-ionic surfactant labeled for aquatics should be added to the spray mixture. Follow all directions and use precautions on the label of the surfactant. Thorough wetting of the foliage and stems is necessary to achieve satisfactory control. A minimum spray volume of 50 gallons per acre is recommended for ground broadcast applications.

If a backpack sprayer is used, a spray mixture containing 1% to 1.5% of Renovate 3 or 5 to 7.6 fl oz of Renovate 3 per 4 gallons of water should be used. All phragmites foliage should be thoroughly wetted.

Aerial application by helicopter may be needed when treating restoration sites that are inaccessible, remote, difficult to traverse, isolated, or otherwise unsuited to ground application, or in circumstances where invasive exotic weeds dominate native plant populations over extensive areas and efforts to restore native plant diversity are being conducted. By air, apply in a minimum spray volume of 30 gallons per acre using Thru-Valve or Microfoil boom only.

- **Recreational Use of Water in Treatment Area:** There are no restrictions on use of water in the treatment area for recreational purposes, including swimming and fishing.
- **Livestock Use of Water from Treatment Area:** There are no restrictions on livestock consumption of water from the treatment area.

TERRESTRIAL SITES ASSOCIATED WITH WETLAND AREAS

- Apply no more than 2 lb ae of triclopyr (2/3 gallon of Renovate 3) per acre per growing season on range and pasture sites, including rights-of-way, fence rows or any area where grazing or harvesting is allowed.
- On forestry sites, Renovate 3 may be used at rates up to 6 lb ae of triclopyr (2 gallons of Renovate 3) per acre per year.

Use Renovate 3 at rates of 3/4 to 6 lb ae of triclopyr (1/4 to 2 gallons of Renovate 3) per acre to control broadleaf weeds and woody plants. In all cases use the amount specified in enough water to give uniform and complete coverage of the plants to be controlled. Use only water suitable for spraying. Use a labeled non-ionic surfactant for all foliar applications. When using surfactants, follow the use directions and precautions listed on the surfactant manufacturer's label. Use the higher recommended concentrations of surfactant in the spray mixture when applying lower spray volumes per acre. The order of addition to the spray tank is water, spray thickening agent (if used), additional herbicide (if used), and Renovate 3. A labeled aquatic surfactant should be added to the spray tank last or as recommended on the product label. If combined with emulsifiable concentrate herbicides, moderate continuous adequate agitation is required.

Before using any recommended tank mixtures, read the directions and all use precautions on both labels.

For best results, apply when woody plants and weeds are actively growing. When hard to control species such as ash, blackgum, choke cherry, maples, or oaks are prevalent and during applications made in late summer when the plants are mature and during drought conditions, use the higher rates of Renovate 3.

When using Renovate 3 in combination with a 2,4-D herbicide approved for aquatic use, such as DMA 4 IVM, generally the higher rates should be used for satisfactory brush control.

Use the higher dosage rates when brush approaches an average of 15 feet in height or when the brush covers more than 60% of the area to be treated. If lower rates are used on hard to control species, re-sprouting may occur the year following treatment.

High Volume Foliage Treatment

For control of woody plants, use Renovate 3 at the rate of 3 to 6 lb ae of triclopyr (1 to 2 gallons of Renovate 3) per 100 gallons of spray solution, or Renovate 3 at 3/4 to 3 lb ae of triclopyr (1 to 4 quarts of Renovate 3) may be tank mixed with 1/4 to 1/2 gallons of 2,4-D 3.8 lb amine, like DMA 4 IVM, diluted to make 100 gallons of spray solution. Apply at a volume of 100 to 400 gallons of total spray per acre depending upon size and density of woody plants. Coverage should be thorough to wet all leaves, stems, and root collars. (See *General Use Precautions and Restrictions*.) Do not exceed the maximum allowable use rate of 6 lb ae of triclopyr (2 gallons of Renovate 3) per acre per growing season.

Low Volume Foliage Treatment

To control susceptible woody plants, apply up to 15 lb ae of triclopyr (5 gallons of Renovate 3) in 10 to 100 gallons of finished spray. The spray concentration of Renovate 3 and total spray volume per acre may be adjusted according to the size and density of target woody plants and kind of spray equipment used. With low volume sprays, use sufficient spray volume to obtain uniform coverage of target plants including the surfaces of all foliage, stems, and root collars (see *General Use Precautions and Restrictions*). For best results, a labeled aquatic surfactant should be added to all spray mixtures. Match equipment and delivery rate of spray nozzles to height and density of woody plants. When treating tall, dense brush, a truck mounted spray gun with spray tips that deliver up to 2 gallons per minute at 40 to 60 psi may be required. Backpack or other types of specialized spray equipment with spray tips that deliver less than 1 gallon of spray per minute may be appropriate for short, low to moderate density brush.

Cut Surface Treatments (Woody Plants)

Individual plant treatments such as basal bark and cut surface applications may be used on any use site listed on this label at a maximum use rate of 2.67 gallons of Renovate 3 (8 lb ae of triclopyr) per acre. These types of applications are made directly to ungrazed parts of plants and, therefore, are not restricted by the grazing maximum rate of 2/3 of a gallon of Renovate 3 (2 lb ae of triclopyr) per acre.

To control unwanted trees and other listed woody plants, apply Renovate 3, either undiluted or diluted in a 1 to 1 ratio with water as directed below.

With Tree Injector Method

Apply by injecting 1/2 milliliter of undiluted Renovate 3 or 1 milliliter of the diluted solution through the bark at intervals of 3 to 4 inches between centers of the injector wound. The injections should completely surround the tree at any convenient height. **Note: No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is injected directly into plants.**

With Hack and Squirt Method

Make cuts at a convenient height around the tree trunk with a hatchet or similar equipment so that the cuts overlap slightly and make a continuous circle around the trunk. Spray 1/2 milliliter of undiluted Renovate 3 or 1 milliliter of the diluted solution into each cut.

With Frill or Girdle Method

Make a single girdle through the bark completely around the tree at a convenient height. Wet the cut surface with undiluted or diluted solution.

Both of the above methods may be used successfully at any season except during periods of heavy sap flow of certain species - for example, maples.

Stump Treatment

Spray or paint the cut surfaces of freshly cut stumps and stubs with undiluted Renovate 3. The cambium area next to the bark is the most vital area to wet.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. Open dumping is prohibited.

Pesticide Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Nonrefillable containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Nonrefillable containers 5 gallons or larger:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or larger:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

TERMS AND CONDITIONS OF USE

If terms of the following *Warranty Disclaimer*, *Inherent Risks of Use*, and *Limitation of Remedies* are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under *Warranty Disclaimer*, *Inherent Risks of Use* and *Limitations of Remedies*.

WARRANTY DISCLAIMER

SePRO Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation or the seller. All such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such loss or damage in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer*, *Inherent Risks of Use*, and this *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer* or this *Limitation of Remedies* in any manner.

EPA Stamped Notification: 06/13/2008

© Copyright 2012 SePRO Corporation

® Renovate is a registered trademark of Dow AgroSciences LLC



SePRO Corporation
11550 North Meridian Street, Suite 600
Carmel, IN 46032, U.S.A.