

**DATE:** 4.3.20  
**FORM:** 1791-G  
**NAME:** Clipper Herbicide  
**KIND:** Booklet Label  
**AMOUNT:** 5 Pounds  
**ACCEPTED**

Classified for  
 "RESTRICTED USE"  
 in New York State  
 under 6NYCRR Part 326

FOR REGISTRATION

FLUMIOXAZIN GROUP 14 HERBICIDE

JUNE 23 2020


New York State Department  
 of Environmental Conservation  
 Division of Materials Management  
 Pesticide Product Registration



Doc ID: 568215

# Clipper®

## HERBICIDE



**FOR THE MANAGEMENT OF UNDESIRABLE AQUATIC  
 VEGETATION IN SLOW MOVING OR QUIESCENT WATERS**

Active Ingredient	By Wt
* Flumioxazin .....	51%
Other Ingredients .....	49%
Total	100%

\* 2-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione  
 Clipper® Herbicide is a water dispersible granule containing 51% active ingredient.

EPA Reg. No. 59639-161 EPA Est. 1773-IA-1® 39578-TX-1© 47857-CA-1  
 Superscript is first letter of lot number.

**KEEP OUT OF  
 REACH OF CHILDREN  
 CAUTION**  
 SEE NEXT PAGE FOR ADDITIONAL  
 PRECAUTIONARY STATEMENTS

**NET WEIGHT 5 POUNDS**



**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS & DOMESTIC ANIMALS**  
**CAUTION**

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing dust and spray mist. Avoid contact with skin, eyes or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or the toilet.

**FIRST AID**

- If inhaled:** Move person to fresh air.  
If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.  
Call a poison control center or doctor for further treatment advice.
- If on skin or clothing:** Take off contaminated clothing.  
Rinse skin immediately with plenty of water for 15-20 minutes.  
Call a poison control center or doctor for treatment advice.
- If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes.  
Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.  
Call a poison control center or doctor for treatment advice.
- If swallowed:** Call a poison control center or doctor immediately for treatment advice.  
Have person sip a glass of water if able to swallow.  
Do not induce vomiting unless told to by a poison control center or doctor.  
Do not give anything by mouth to an unconscious person.

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some of the materials that are chemical-resistant to this product are listed below.

**Applicators and other handlers must wear:**

- long-sleeved shirt
- long pants

- chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- shoes and socks.

**User Safety Requirements**

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS:**

If not used in accordance with directions on the label, this product can be toxic to non-target plants and aquatic invertebrates. Do not apply to water except as specified on the label. Drift and runoff may be hazardous to non-target plants and aquatic organisms in neighboring areas, if not used in accordance to label directions. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

This pesticide is toxic to plants and should be used strictly in accordance with the drift and runoff precautions on this label in order to minimize off-site exposures.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the treatment area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**DISCLAIMER, RISKS OF USING THIS PRODUCT,  
LIMITED WARRANTY  
AND LIMITATION OF LIABILITY**

**IMPORTANT:** Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

**RISKS OF USING THIS PRODUCT**

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

**LIMITED WARRANTY**

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law AND AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED. No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

**LIMITATION OF LIABILITY**

To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect

(continued)

(continued)

or special damages resulting from the use or handling of this product. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

**PROMPT NOTICE OF CLAIM**

To the extent consistent with applicable law allowing such requirements Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from the date of application so that an immediate inspection of the affected property can be made.

To the extent consistent with applicable law if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

**NO AMENDMENTS**

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing **Disclaimer, Risks of Using This Product, Limited Warranty** and **Limitation of Liability**, which may not be modified by any oral or written agreement.

**TANK MIXES**

**NOTICE:** Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

## PRODUCT INFORMATION

*Clipper* Herbicide is a fast acting contact herbicide that controls selected submersed, emergent and floating aquatic weeds. It is most effective when applied to young, actively growing weeds in water with a pH of less than 8.5.

*Clipper* Herbicide may be applied to the following quiescent or slow moving bodies of water:

- Bayous
- Canals
- Drainage ditches
- Lakes
- Marshes
- Ponds (including golf course ponds)
- Reservoirs

Application of *Clipper* Herbicide to public aquatic areas may require special approval and/or permits. Consult with local state agencies, if required.

## IRRIGATION RESTRICTIONS FOLLOWING APPLICATION

Application Method	Application Rate	Average Water Depth	Turf and Landscape Ornamentals	Ornamentals Grown for Production in Greenhouse and Nursery
Surface Spray	6 to 12 oz per surface acre	Greater than 3 feet	None	5 days
		Less than 3 feet	12 hours	5 days
Subsurface	Less than 200 ppb	N/A	1 day	5 days
	200 to 300 ppb	N/A	2 days	5 days
	300 to 400 ppb	N/A	3 days	5 days

### Weed Resistance Management

For resistance management, please note that *Clipper* Herbicide contains a Group 14 herbicide. Any weed population may contain plants naturally resistant to *Clipper* Herbicide and other Group 14 herbicides. The resistant individuals may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of *Clipper* Herbicide or other Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.

(continued)

## RESTRICTIONS

- Do not exceed 400 ppb of *Clipper* Herbicide during any one application.
- Do not re-treat the same section of water with *Clipper* Herbicide more than 6 times per year.
- Do not apply to intertidal or estuarine areas.
- Do not retreat the same section of water within 28 days of application. In areas with dense weed vegetation only treat 1/2 the water body at one time and wait 10-14 days before treating the remaining area.
- Do not use in water utilized for crawfish farming.
- **Do not use treated water for irrigation purposes on food crops until at least five (5) days after application.**

### Notes

- Treated water may be used for irrigation purposes on turf and landscape ornamentals as outlined in the *Irrigation Restrictions Following Application* table.
- There is no post-application holding restriction against use of treated water for drinking or recreational purposes (e.g., swimming, fishing).

(continued)

- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and other management practices.

(continued)

(continued)

- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance management and/or integrated weed-management strategies for specific weed biotypes.
- For further information or to report suspected resistance, contact Valent U.S.A. LLC at 800-682-5368.

## SPRAY DRIFT MANAGEMENT FOR FOLIAR OR SURFACE APPLICATIONS

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

Do not spray *Clipper* Herbicide under circumstances where spray droplets may drift on to unprotected persons, or plantings of food, forage or crops that might be damaged, or rendered unfit for sale, use or consumption. These precautions are not applicable for subsurface injection by closed systems.

- Use the largest droplet size consistent with acceptable efficacy. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. For ground boom and aerial applications, use medium or coarser spray nozzles according to ASAE 572 definition for standard nozzles or a volume mean diameter (VMD) of 300 microns or greater for spinning atomizer nozzles.

- Make aerial, ground or watercraft-based surface applications when wind velocity favors on-target product deposition. Apply only when the wind speed is less than or equal to 10 mph.
- Do not make aerial or ground applications into areas of temperature inversions. Inversions are characterized by stable air and increasing temperatures with increasing distance above the ground. Mist or fog may indicate the presence of an inversion in humid areas. Where permissible by local regulations, the applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.
- Low humidity and high temperatures increase the evaporation rate of spray droplets, and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperatures.

Properly maintain and calibrate all aerial, ground and water based application equipment.

Where states have more stringent regulations, they should be observed.

## APPLICATION AND SPRAYER INFORMATION

### Mixing Instructions

- Fill clean spray tank 1/2 full of desired level with water and add buffering agent if necessary.
- Add the required amount of *Clipper* Herbicide to the spray tank while agitating.
- Fill spray tank to desired level with water. Ensure that *Clipper* Herbicide is thoroughly mixed before making applications. Agitation should continue until spray solution has been applied.
- Mix only the amount of spray solution that can be applied the day of mixing. Apply *Clipper* Herbicide within 12 hours of mixing.

### ADDITIVES

When applying *Clipper* Herbicide to the foliage of floating or emerged aquatic weeds, mix with an adjuvant approved for use in aquatic sites. Valent recommends the use of a Chemical Producers and Distributors Association certified adjuvant. Mix *Clipper* Herbicide with a non-ionic surfactant containing at least 80% active ingredient. Follow adjuvant manufacturer's label rates. Mixing compatibility should be verified by a jar test before using.

### Jar Test to Determine Compatibility of Adjuvants and *Clipper* Herbicide

Conduct a jar test before mixing commercial quantities of *Clipper* Herbicide, when using for the first time, when using new adjuvants or when a new water source is being used.

1. Add 1 pt of water to a quart jar. The water should be from the same source and have the same temperature as the water used in the spray tank mixing operation.
2. Add 3 grams (approximately 1 level tsp) of *Clipper* Herbicide for the 8 oz/A rate or 4 grams (approximately 1-1/2 tsp) for 12 oz/A rate to the jar. Gently mix until product disperses.
3. Add 60 ml (4 Tbsp or 2 fl oz) of additive to the quart jar and gently mix.
4. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
5. An ideal tank mix combination will be uniform and free of suspended particles. If any of the following conditions are observed the choice of adjuvant should be questioned:
  - a) Layer of oil or globules on the solution surface.
  - b) Flocculation: Fine particles in suspension or as a layer on the bottom of the jar.
  - c) Clabbering: Thickening texture (coagulated) like gelatin.

### Sprayer Cleanup

If spray equipment is dedicated to application of aquatic herbicides, the following steps are recommended to clean the spray equipment:

- Completely drain the spray tank and rinse the application equipment thoroughly, including the inside and outside of the tank and all in-line screens.

If spray equipment will be used for purposes other than applying aquatic herbicides, it must be thoroughly cleaned following application of *Clipper* Herbicide. The following steps must be used to clean the spray equipment:

1. Completely drain the spray tank and rinse the application equipment thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the tank with clean water and flush all hoses, booms, screens and nozzles.
3. Top off tank with clean water.
4. Circulate through sprayer for 5 minutes.
5. Then flush all hoses, booms, screens and nozzles for a minimum of 15 minutes.
6. Drain tank completely.
7. Remove all nozzles and screens and rinse them with clean water.

### AERIAL APPLICATION

To obtain satisfactory weed control, aerial application of *Clipper* Herbicide, must provide uniform coverage of surface weeds and sufficient contact time. When applied by air, *Clipper* Herbicide may not provide adequate control of some submersed weeds. Do not apply

by air when significant drift on to non-target plants may occur or when wind velocity is more than 10 mph. Avoid spraying *Clipper* Herbicide within 200 feet of dwellings, adjacent sensitive crops or environmentally sensitive areas. To obtain satisfactory application and avoid drift, the following directions must be observed:

#### Volume and Pressure

Apply *Clipper* Herbicide in a minimum of 5 gallons of water per acre, with a maximum spray pressure of 40 PSI. Application at less than 5 gallons per acre may not provide adequate weed control. Higher gallonage applications generally provide more consistent weed control.

#### Nozzles and Nozzle Operation

Use nozzles that produce flat or hollow cone spray patterns. Use non-drip type nozzles such as diaphragm type nozzles to avoid unwanted discharge of spray solution. The nozzle must be directed toward the rear of the aircraft, at an angle between 0° and 15° downward. Do not place nozzles on the outer 25% of the wings or rotors.

#### Adjuvants

Refer to the additive section or the tank mix partners label for adjuvant recommendation.

### DIRECTIONS FOR USE TO CONTROL FLOATING AND EMERGED WEEDS USING SURFACE APPLICATION

*Clipper* Herbicide will control weeds and algae listed in Table 1, Floating and Emerged Weeds when applied as a broadcast spray with appropriate equipment. For best results, apply *Clipper* Herbicide to the foliage of actively growing weeds.

**Table 1. Floating and Emerged Weeds**

Common Name	Scientific Name
Alligator Weed	<i>Alternanthera philoxeroides</i>
Duckweed	<i>Lemna</i> spp.
Frog's-bit	<i>Limnobium spongia</i>
Water Fern	<i>Salvinia</i> spp.
Water Lettuce	<i>Pistia stratiotes</i>
Watermeal	<i>Wolffia</i> spp.
Water Pennywort*	<i>Hydrocotyle</i> spp.
Filamentous algae*	<i>Pithophora</i>
Filamentous algae*	<i>Cladophora</i>

\*Not for use in California.

#### Surface Application

Apply *Clipper* Herbicide as a broadcast spray at 6 to 12 ounces of formulated product per acre plus an adjuvant approved for use in aquatics.

*Clipper* Herbicide is a contact herbicide that quickly degrades in the water column so plants that do not initially come in contact with the herbicide will not be controlled. Apply *Clipper* Herbicide in a minimum of 30 gallons of water per acre to all areas of the water body where weeds exist. Coverage is essential for effective control as all floating weeds need to be exposed to lethal concentrations in all parts of the water body. Any untreated escapes or re-introductions of plants that were not treated will reestablish in areas where surface weeds had previously been controlled. If a second application is required to provide control, it is recommended that a treatment be made once the return of these weeds is first observed, but no sooner than 28 days after the last treatment.

Application of *Clipper* Herbicide during early morning hours may enhance weed control. When applying to densely packed actively growing surface weeds, ensure adequate coverage. Rapid decomposition of vegetation resulting from herbicide treatment can result in loss of oxygen in water. A sudden decrease in dissolved oxygen can result in fish suffocation. If aquatic vegetation is dense, treat floating surface weeds in sections to avoid a rapid decrease in dissolved oxygen.

*Clipper* Herbicide may be tank mixed with 2,4-D, diquat, glyphosate or other registered foliar applied herbicides for enhanced control of floating and emergent weeds.

Consult a manufacturer's label for specific rate restrictions and weeds controlled. Always follow the most restrictive label restrictions and precautions for all products used when making an application involving tank mixes.

### Application Equipment

Apply *Clipper* Herbicide with sprayers equipped with nozzles designed to deliver the desired spray pressure and spray volume. Apply by backpack or handgun sprayer, airboat, helicopter, airplane or other application equipment that will ensure thorough coverage of target plant foliage.

## DIRECTIONS FOR USE TO CONTROL SUBMERSED AND FLOATING WEEDS USING SUBSURFACE APPLICATION

*Clipper* Herbicide will control submersed and floating weeds listed in Table 2, Submersed and Floating Weeds Controlled by Subsurface Application, when applied subsurface with appropriate equipment.

**Table 2. Submersed and Floating Weeds Controlled by Subsurface Application**

Common Name	Scientific Name
Coontail	<i>Ceratophyllum demersum</i>
Duckweed	<i>Lemna</i> spp.
Fanwort	<i>Cabomba caroliniana</i>
Hydrilla	<i>Hydrilla verticillata</i>
Hygrophila	<i>Hygrophila polysperma</i>
Naiad, Southern	<i>Najas guadalupensis</i>
Pondweed, Curlyleaf*	<i>Potamogeton crispus</i>
Pondweed, Sago*	<i>Potamogeton pectinatus</i>
Pondweed, Variable-Leaf*	<i>Potamogeton diversifolius</i>
Water Fern	<i>Salvinia</i> spp.
Water Lettuce	<i>Pistia stratiotes</i>
Watermeal	<i>Wolffia</i> spp.
Watermilfoil, Eurasian	<i>Myriophyllum spicatum</i>
Watermilfoil, Variable-Leaf	<i>Myriophyllum heterophyllum</i>

\*Not for use in California.

### Subsurface Application

Apply *Clipper* Herbicide at a rate that will produce an initial concentration of 200 to 400 ppb (of active ingredient flumioxazin) in the water column.

*Clipper* Herbicide is rapidly absorbed by target plants, but also breaks down quickly in water with a pH greater than 8.5. The pH of water surrounding mats of submersed vegetation can exceed 8.5 by early to mid-day, due to photosynthetic processes. Application of *Clipper* Herbicide under these conditions may provide only partial weed control, and regrowth is likely. For best control, apply *Clipper* Herbicide in a minimum of 30 gallons of water per acre in the early morning to actively growing weeds and early in the season before surface matting occurs. Complete coverage and sufficient contact time of submersed weeds with *Clipper* Herbicide is required for optimal performance. Application of *Clipper* Herbicide with subsurface trailing hoses designed to distribute the herbicide within the plant stand will provide more effective and longer term control of submersed weeds. Use Table 3, Subsurface Application Rates to determine the amount of *Clipper* Herbicide needed to achieve desired concentration at different water depths. Use higher concentrations when weed biomass is heavy and/or weeds are more mature and topped out. Any untreated plants that are left in the water column can re-infest treated areas that had previously been controlled. If a second application is required to provide control, it is recommended that a treatment be made once the return of these weeds is

first observed, but no sooner than 28 days after the last treatment.

When applying *Clipper* Herbicide to densely packed actively growing submersed weeds, a rapid decomposition of vegetation resulting from herbicide treatment can result in loss of oxygen in water. A sudden decrease in dissolved oxygen can result in fish suffocation. If aquatic vegetation is dense, treat submersed weeds in sections to avoid a rapid decrease in dissolved oxygen.

*Clipper* Herbicide may be tank mixed with other registered submersed applied herbicides for enhanced control of submersed and floating weeds.

### Application Equipment

To improve distribution in the water column and ensure adequate coverage, apply *Clipper* Herbicide with subsurface trailing hoses in order to place the herbicide under the surface and throughout the biomass of aquatic vegetation. Keep swath width to a minimum in order to maximize contact with submersed aquatic vegetation.

### Information on Hydrilla Control in Florida

*Clipper* Herbicide should be applied as a subsurface treatment for hydrilla control. For best control of hydrilla apply during the late Winter/early Spring and/or early to late Fall. Efficacy of *Clipper* Herbicide will be enhanced at these timings due to lower potential biomass present and lower pH of the water. If applied to mature topped out hydrilla, *Clipper* Herbicide will cause some discoloration and loss of growing tips, but regrowth will be rapid.

Tank mixing *Clipper* Herbicide with other registered herbicides is recommended, especially if hydrilla is approaching maturity or biomass is heavy.

**Table 3. Subsurface Application Rates**

Water Depth (feet)	Pounds of <i>Clipper</i> Herbicide required per surface acre to achieve desired water concentration		
	200 ppb	300 ppb	400 ppb
1	1.1	1.6	2.1
2	2.1	3.2	4.2
3	3.2	4.8	6.4
4	4.2	6.4	8.5
5	5.3	8.0	10.6
6	6.4	9.5	12.7
7	7.4	11.1	14.8

Example: to achieve an initial concentration of 200 ppb of flumioxazin in a 4 foot deep water column, apply 4.2 lb of *Clipper* Herbicide per surface acre.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

### PESTICIDE STORAGE

Keep pesticide in original container.

Store in a cool, dry, secure place.

Do not put formulation or dilute spray solution into food or drink containers.

Do not contaminate food or foodstuffs.

Do not store or transport near feed or food.

Not for use or storage in or around the home.

For help with any spill, leak, fire or exposure involving this material, call day or night **(800) 892-0099**.

### PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

©2020 Valent U.S.A. LLC

*Clipper* is a registered trademark of Valent U.S.A. LLC

Manufactured for:

**Valent U.S.A. LLC**

P.O. Box 8025

Walnut Creek CA 94596-8025

Made in U.S.A.

Form 1791-G

EPA Reg. No. 59639-161

EPA Reg. No. 59639-161 EPA Est. 1773-1A-1®  
39578-TX-1© 47857-CA-1

Superscript is first letter of lot number.


059639-00161.20180612.CL.PAMEND.FINAL  
SAL20180614





# Clipper®

## HERBICIDE



**FOR THE MANAGEMENT OF UNDESIRABLE AQUATIC  
VEGETATION IN SLOW MOVING OR QUIESCENT WATERS**

Active Ingredient	By Wt
* Flumioxazin .....	51%
Other Ingredients .....	49%
Total	100%

\* 2-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione

Clipper® Herbicide is a water dispersible granule containing 51% active ingredient.

EPA Reg. No. 59639-161 EPA Est. 1773-IA-1<sup>®</sup>, 39578-TX-1<sup>®</sup>, 47857-CA-1  
Superscript is first letter of lot number.

**KEEP OUT OF  
REACH OF CHILDREN  
CAUTION**  
SEE INSIDE BOOKLET FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS

**NET WEIGHT 5 POUNDS**